Enhancements in the Evaluation of Plutonium Deposited in Lung Tissue

David P. Hickman and <u>Sandi Fisher</u> Lawrence Livermore National Laboratory

The in vivo measurement of plutonium has traditionally relied upon the measurement of a particle-associated radionuclide such as ²⁴¹Am or by directly measuring the x-rays emitted from the decay of the plutonium isotopes. Other methods of directly measuring plutonium in vivo have also used the low abundance gamma emissions from plutonium. All of the methods used for the in vivo measurement of Plutonium are highly dependent on the transmission of the photons through interfering structures such as muscle and bone. Recent studies at Lawrence Livermore National Laboratory have identified the optimum method as a function of chest wall thickness for measuring pure ²³⁸Pu and ²³⁹Pu deposited in the lung.

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.