


Ultrafast measurements explain quantum dot voltage drop

March 22, 2017



Insights pave way for solar cells and photodetectors based on tunable nanoparticles

LOS ALAMOS, N.M., March 22, 2017—Solar cells and photodetectors could soon be made from new types of materials based on semiconductor quantum dots, thanks to new insights based on ultrafast measurements capturing real-time photoconversion processes.

“Our latest ultrafast electro-optical spectroscopy studies provide unprecedented insights into the photophysics of quantum dots,” said lead researcher Victor Klimov, a physicist specializing in semiconductor nanocrystals at Los Alamos National Laboratory, “and this new information helps perfect the materials’ properties for applications in practical photoconversion devices. Our new experimental technique allows us to follow a chain of

events launched by femtosecond laser pulses and pin down processes responsible for efficiency losses during transformation of incident light into electrical current.”

Photoconversion is a process wherein the energy of a photon, or quantum of light, is converted into other forms of energy, for example, chemical or electrical. Semiconductor quantum dots are chemically synthesized crystalline nanoparticles that have been studied for more than three decades in the context of various photoconversion schemes including photovoltaics (generation of photo-electricity) and photo-catalysis (generation of “solar fuels”). The appeal of quantum dots comes from the unmatched tunability of their physical properties, which can be adjusted by controlling the size, shape and composition of the dots.

2:30

Controlling the properties of matter with quantum dots

At Los Alamos, the research connects to the institutional mission of solving national security challenges through scientific excellence, in this case focusing on novel physical principles for highly efficient photoconversion, charge manipulation in exploratory device structures and novel nanomaterials.

The interest in quantum dots as solar-cell materials has been motivated by their tunable optical spectra as well as interesting new physics such as [high-efficiency carrier multiplication](#), that is, generation of multiple electron-hole pairs by single photons. This effect, discovered by Los Alamos researchers in 2004, resulted in the surge of activities in the area of quantum dot solar cells that quickly pushed the efficiencies of practical devices to more than 10 percent.

Further progress in this area has been hindered by the challenge of understanding the mechanisms of electrical conductance in quantum dot solids and the processes that limit the charge transport distance. One specific and persistent challenge of great importance from the standpoint of photovoltaic (PV) applications, Klimov said, is understanding the reasons underlying a considerable loss in photovoltage compared to predicted theoretical limits—a problem with quantum dot solar cells known as a “photovoltage deficit.” Los Alamos researchers at the Center for Advanced Solar Photophysics (CASP) helps answer some of the above questions.

By applying a combination of ultrafast optical and electrical techniques, the Los Alamos scientists have been able to resolve step-by-step a sequence of events involved in photoconversion in quantum dot films from generation of an exciton to electron-hole separation, dot-to-dot charge migration and finally recombination.

The high temporal resolution of these measurements (better than one billionth of a second) enabled the team to reveal the cause of a large drop of the electron energy, which results from very fast electron trapping by defect-related states. In the case of practical devices, this process would result in reduced photovoltage. The newly conducted studies establish the exact time scale of this problematic trapping process and suggest that a moderate (less than ten-fold) improvement in the electron mobility should allow for collecting photogenerated charge carriers prior to their relaxation into lower-energy states. This would produce a dramatic boost in the photovoltage and therefore increase the overall device efficiency.

Another interesting effect revealed by these studies is the influence of electron and hole “spins” on photoconductance. Usually spin properties of particles (they can be thought of as the rate and direction of particle rotation around its axis) are invoked in the

case of interactions with a magnetic field. However, previously it was found that even a weak interaction between spins of an electron and a hole (so-called "spin-exchange" interaction) has a dramatic effect on light emission from the quantum dots.

The present measurements reveal that these interactions also affect the process of electron-hole separation between adjacent dots in quantum-dot solids. Specifically these studies suggest that future efforts on high-sensitivity quantum-dot photodetectors should take into consideration the effect of exchange blockade, which otherwise might inhibit low-temperature photoconductance.

Quantum dot materials have been at the heart of research at the Los Alamos [Center for Advanced Solar Photophysics](#), which has investigated their application to solar-energy technologies such as luminescent sunlight collectors for solar windows and low-cost PV cells processed from quantum dot solutions.

The paper, in this week's Nature Physics Journal: "[Electron-hole exchange blockade and memory-less recombination in photoexcited films of colloidal quantum dots](#)," [Andrew F. Fidler](#), [Jianbo Gao](#) & [Victor I. Klimov](#)

Funding: This work was funded by the Department of Energy's Office of Science through a grant to the Center for Advance Solar Photophysics (CASP), a DOE Energy Frontier Research Center.

Los Alamos National Laboratory

www.lanl.gov

(505) 667-7000

Los Alamos, NM

Operated by Los Alamos National Security, LLC for the Department of Energy's NNSA

