

# Higgs Bosons in Left-Right Supersymmetric Models



K.S. Babu  
Oklahoma State University

Santa Fe 2014 Summer Workshop  
“LHC After the Higgs”  
June 30 – July 4, 2014

# Outline

- ❧ Why Left-Right Supersymmetry?
- ❧ Higgs Boson spectrum in Supersymmetric Left-Right models
  - Models with Higgs triplets + bidoublets
  - Models with Higgs doublets + bidoublets: Inverse seesaw models
  - Models with Higgs doublets: Universal seesaw models
  - $E_6$  motivated Left-Right models
- ❧ Emergence of a Light Doubly Charged Higgs boson and associated phenomenology
- ❧ Conclusions

Based on: K.S. Babu and Ayon Patra, 2014 (to appear);  
K.S. Babu, Ayon Patra, Santsh Kumar Rai, Phys. Rev. D88, 055006 (2013).

# Highlights

- Lightest neutral Higgs boson mass can easily be 125 GeV with low  $\tan \beta \sim 1$ , with light stops of mass below 1 TeV and negligible stop mixing. This results in reduced fine-tuning.
- Non-decoupling  $SU(2)_R$  D-term makes this possible (within MSSM,  $\tan \beta > 5$  and stop mass  $> 1$  TeV is needed).
- A light doubly charged Higgs boson is predicted in models with Higgs triplets, which acquires its mass entirely from radiative corrections.
- Several variants supersymmetrized for the first time.

# Why Left-Right Supersymmetry?

Addresses some important open questions of the Standard Model:

- Understanding the origin of parity violation.
- Natural setting for small neutrino mass via seesaw mechanism.
- Provides a simple solution to the Strong CP Problem.
- SUSY stabilizes the Higgs boson mass.
- Natural dark matter candidate without assuming R-parity.
- Solves the SUSY CP problem.

J. C. Pati and A. Salam, Phys. Rev. D (1973) ;

R. N. Mohapatra and J. C. Pati, Phys. Rev. D (1975) ;

G. Senjanovic and R. N. Mohapatra, Phys. Rev. D (1975).

# How Left-Right Symmetry solves the Strong CP Problem & SUSY CP Problem

Strong interaction sector admits a term that violates both CP and P:

$$\mathcal{L}_{QCD} = \frac{\theta g^2}{32\pi^2} G_{\mu\nu}^a \tilde{G}^{a\mu\nu}$$

Leads to the physical observable

$$\bar{\theta} = \theta + \text{Arg}(\text{Det } M_q) \quad M_q \rightarrow \text{quark mass matrix}$$

Neutron electric dipole moment limits imply  $\bar{\theta} < 10^{-10}$  .

With left-right symmetry,  $\theta = 0$  due to Parity;  $M_q$  is Hermitian also due to Parity, and thus  $\bar{\theta} = 0$  at tree level.

Induced  $\bar{\theta}$  is small and consistent.

Parity symmetry also makes the new SUSY phases zero, thus solving the SUSY CP problem.

R.N. Mohapatra and A. Rasin, Phys. Rev. Lett. 76, 3490 (1996);

R. Kuchimanchi, Phys. Rev. Lett. 76, 3486 (1996);

K.S. Babu, B. Dutta, R.N. Mohapatra, Phys. Rev. D60, 095004 (1999);

K.S. Babu, B. Dutta, R.N. Mohapatra, Phys. Rev. D65, 016005 (2002).

# Higgs Boson masses in Supersymmetric Left-Right models

- ∞ Supersymmetric versions of left-right symmetry has the gauge group extended to  $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ .
- ∞ Electric charge defined as  $Q = I_{3L} + I_{3R} + \frac{B-L}{2}$ .
- ∞ The  $SU(2)_R \times U(1)_{B-L}$  symmetry is broken spontaneously to  $U(1)_Y$  at a high scale.
- ∞ Here we consider several variations with different symmetry breaking sectors. Either Higgs triplets or Higgs doublets are used for the high scale symmetry breaking.
- ∞ Higgs boson spectrum is calculated in each case.

# Fermion Content of Left-Right Models

∞ The common Quark and Lepton sectors of all models are:

$$Q(3,2,1,1/3) = \begin{bmatrix} u \\ d \end{bmatrix} \quad ; \quad Q^c(3^*,1,2,-1/3) = \begin{bmatrix} d^c \\ -u^c \end{bmatrix}$$

$$L(1,2,1,-1) = \begin{bmatrix} \nu_e \\ e \end{bmatrix} \quad ; \quad L^c(1,1,2,1) = \begin{bmatrix} e^c \\ -\nu_e^c \end{bmatrix}$$

- The Right-handed quarks and leptons are doublets of  $SU(2)_R$ .
- Presence of Right-handed neutrino required by gauge structure.

∞ Each model has to explain

- Consistent symmetry breaking mechanism.
- Quarks and Lepton masses and CKM mixing.
- Small neutrino mass generation.

# Models with Higgs triplets + bidoublets

Most straightforward way to break symmetry and generate large mass for right-handed neutrinos.

Under  $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$

$$\Delta^c(1,1,3,-2) = \begin{bmatrix} \frac{\delta^{c-}}{\sqrt{2}} & \delta^{c0} \\ \delta^{c--} & -\frac{\delta^{c-}}{\sqrt{2}} \end{bmatrix} \quad \bar{\Delta}^c(1,1,3,2) = \begin{bmatrix} \frac{\bar{\delta}^{c+}}{\sqrt{2}} & \bar{\delta}^{c++} \\ \bar{\delta}^{c0} & -\frac{\bar{\delta}^{c+}}{\sqrt{2}} \end{bmatrix}$$

$$\Delta(1,3,1,2) = \begin{bmatrix} \frac{\delta^+}{\sqrt{2}} & \delta^{++} \\ \delta^0 & -\frac{\delta^+}{\sqrt{2}} \end{bmatrix} \quad \bar{\Delta}(1,3,1,-2) = \begin{bmatrix} \frac{\bar{\delta}^-}{\sqrt{2}} & \bar{\delta}^0 \\ \bar{\delta}^{--} & -\frac{\bar{\delta}^-}{\sqrt{2}} \end{bmatrix}$$

$$\Phi_a(1,2,2,0) = \begin{bmatrix} \phi_1^+ & \phi_2^0 \\ \phi_1^0 & \phi_2^- \end{bmatrix}_a \quad (a=1,2) \quad S(1,1,1,0)$$

R. Kuchimanchi, R. N. Mohapatra, Phys. Rev. D48, 4352 (1993) .

C.S. Aulakh, K. Benakli, G. Senjanovic, Phys. Rev. Lett. 79, 2188 (1997);

C.S. Aulakh, A. Melfo, G. Senjanovic, Phys. Rev. D57, 4174 (1998).

- The  $SU(2)_R$  Higgs boson triplets ( $\Delta^c$  and  $\overline{\Delta}^c$ ) are needed to break  $SU(2)_R \times U(1)_{B-L} \rightarrow U(1)_Y$  without inducing  $R$ -Parity violating couplings.
- The  $SU(2)_L$  Higgs boson triplets ( $\Delta$  and  $\overline{\Delta}$ ) are their parity partners.
- The two bi-doublet Higgs fields  $\Phi_a$  generate the quark and lepton masses and the CKM mixings.
- The optional singlet field  $S$  makes sure that the right-handed symmetry breaking may occur in the supersymmetric limit.
- The non-zero vacuum expectation values of the fields are

$$\langle \delta^{c0} \rangle = v_R, \quad \langle \overline{\delta}^{c0} \rangle = \overline{v}_R, \quad \langle \phi_{1a}^0 \rangle = v_{u_a}, \quad \langle \phi_{2a}^0 \rangle = v_{d_a}$$

where

$$v_R, \overline{v}_R \gg v_{u_a}, v_{d_a}$$

**Previous Higgs boson mass analysis: Zhang, An, Ji, Mohapatra (2008)**

The Yukawa couplings terms in the superpotential are

$$W = Y_u Q^T \tau_2 \phi_1 \tau_2 Q^c + Y_d Q^T \tau_2 \phi_2 \tau_2 Q^c + Y_\nu L^T \tau_2 \phi_1 \tau_2 L^c + Y_l L^T \tau_2 \phi_2 \tau_2 L^c \\ + i(f^* L^T \tau_2 \Delta L + f L^{cT} \tau_2 \Delta^c L^c)$$

- $\nu_e^c$  is heavy and generates small neutrino mass.

The Higgs boson only superpotential is

$$W_{Higgs} = S \left[ \text{Tr}(\lambda^* \Delta \bar{\Delta} + \lambda \Delta^c \bar{\Delta}^c) + \lambda'_{ab} \text{Tr}(\phi_a^T \tau_2 \phi_b \tau_2) - M_R^2 \right] \\ + \text{Tr} \left[ \mu_1 \Delta \bar{\Delta} + \mu_2 \Delta^c \bar{\Delta}^c \right] + \frac{\mu}{2} \text{Tr}(\phi_a^T \tau_2 \phi_b \tau_2) + \frac{\mu_S}{2} S^2$$

The Superpotential is invariant under parity transformation.

- Yukawa coupling matrices are hermitian.
- $\lambda'_{ab}, \mu, \mu_S$  and  $M_R^2$  are real and  $\mu_1 = \mu_2^*$ .
- If the bi-doublet VEVs are real, it can solve the Strong CP and SUSY CP problem.

# Higgs Potential $V_{Higgs} = V_F + V_D + V_{Soft}$

$$\begin{aligned}
 V_F &= \text{Tr} \left| (\lambda \Delta \bar{\Delta}) + (\lambda^* \Delta^c \bar{\Delta}^c) + \frac{\lambda'}{2} (\Phi^T \tau_2 \Phi \tau_2) - M^2 + \mu_S S \right|^2 + \text{Tr} |\mu \Phi + \lambda' S \Phi|^2 \\
 &+ \text{Tr} \left[ |\mu_1 \Delta + \lambda S \Delta|^2 + |\mu_1 \bar{\Delta} + \lambda S \bar{\Delta}|^2 + |\mu_2 \Delta^c + \lambda^* S \Delta^c|^2 \right. \\
 &\left. + |\mu_2 \bar{\Delta}^c + \lambda^* S \bar{\Delta}^c|^2 \right],
 \end{aligned}$$

$$\begin{aligned}
 V_D &= \frac{g_L^2}{8} \sum_{a=1}^3 \left| \text{Tr} (2\Delta^\dagger \tau_a \Delta + 2\bar{\Delta}^\dagger \tau_a \bar{\Delta} + \Phi^\dagger \tau_a \Phi) \right|^2 \\
 &+ \frac{g_R^2}{8} \sum_{a=1}^3 \left| \text{Tr} (2\Delta^{c\dagger} \tau_a \Delta^c + 2\bar{\Delta}^{c\dagger} \tau_a \bar{\Delta}^c + \Phi^\dagger \tau_a \Phi) \right|^2 \\
 &+ \frac{g_V^2}{2} \left| \text{Tr} (\Delta^\dagger \Delta - \bar{\Delta}^\dagger \bar{\Delta} - \Delta^{c\dagger} \Delta^c + \bar{\Delta}^{c\dagger} \bar{\Delta}^c) \right|^2,
 \end{aligned}$$

$$\begin{aligned}
 V_{Soft} &= m_1^2 \text{Tr} (\Delta^{c\dagger} \Delta^c) + m_2^2 \text{Tr} (\bar{\Delta}^{c\dagger} \bar{\Delta}^c) + m_3^2 \text{Tr} (\Delta^\dagger \Delta) + m_4^2 \text{Tr} (\bar{\Delta}^\dagger \bar{\Delta}) \\
 &+ m_S^2 |S|^2 + m_5^2 \text{Tr} (\Phi^\dagger \Phi) + [\lambda A_\lambda S \text{Tr} (\Delta \bar{\Delta} + \Delta^c \bar{\Delta}^c) + h.c.] \\
 &+ [\lambda' A_{\lambda'} S \text{Tr} (\Phi^T \tau_2 \Phi \tau_2) + h.c.] + (\lambda C_\lambda M^2 S + h.c.) + (\mu_S B_S S^2 + h.c.) \\
 &+ [\mu_1 B_1 \text{Tr} (\Delta \bar{\Delta}) + \mu_2 B_2 \text{Tr} (\Delta^c \bar{\Delta}^c) + \mu B \text{Tr} (\Phi^T \tau_2 \Phi \tau_2) + h.c.].
 \end{aligned}$$

$$M_{11} = \frac{g_L^2(v_1^2 - v_2^2)^2 + g_R^2(v_1^2 - v_2^2)^2 + 8v_1^2v_2^2\lambda'^2}{2(v_1^2 + v_2^2)},$$

$$M_{12} = \frac{v_1v_2(v_1^2 - v_2^2)(g_L^2 + g_R^2 - 2\lambda'^2)}{(v_1^2 + v_2^2)},$$

$$M_{13} = \frac{-g_R^2(v_1^2 - v_2^2)(v_R^2 - \bar{v}_R^2) - 4\lambda\lambda'v_1v_2v_R\bar{v}_R}{\sqrt{(v_1^2 + v_2^2)(v_R^2 + \bar{v}_R^2)}},$$

$$M_{14} = \frac{2[g_R^2(v_1^2 - v_2^2)v_R\bar{v}_R - \lambda\lambda'v_1v_2(v_R^2 - \bar{v}_R^2)]}{\sqrt{(v_1^2 + v_2^2)(v_R^2 + \bar{v}_R^2)}},$$

$$M_{15} = \frac{2\lambda'[-2A_\lambda v_1v_2 + (v_1^2 + v_2^2)(v_S\lambda' + \mu) - \mu_S v_1v_2]}{\sqrt{v_1^2 + v_2^2}},$$

$$M_{22} = \left[ (2g_L^2 + 2g_R^2)v_1^2v_2^2 + 2m_5^2(v_1^2 + v_2^2) + \lambda'^2(v_1^2 - v_2^2)^2 + 2\lambda'^2v_S^2(v_1^2 + v_2^2) \right. \\ \left. + 4\lambda'\mu v_S(v_1^2 + v_2^2) + 2\mu^2(v_1^2 + v_2^2) \right] / (v_1^2 + v_2^2),$$

$$M_{23} = \frac{2[g_R^2v_1v_2(-v_R^2 + \bar{v}_R^2) + \lambda\lambda'(v_1^2 - v_2^2)v_R\bar{v}_R]}{\sqrt{v_1^2 + v_2^2}\sqrt{v_R^2 + \bar{v}_R^2}},$$

$$M_{24} = \frac{-4g_R^2v_1v_2v_R\bar{v}_R - \lambda\lambda'(v_1^2 - v_2^2)(v_R^2 - \bar{v}_R^2)}{\sqrt{v_1^2 + v_2^2}\sqrt{v_R^2 + \bar{v}_R^2}},$$

$$M_{25} = \frac{\lambda'(v_1^2 - v_2^2)(2A_\lambda + \mu_S)}{\sqrt{v_1^2 + v_2^2}},$$

$$M_{33} = \frac{2[(g_R^2 + g_V^2)(v_R^2 - \bar{v}_R^2)^2 + 2\lambda^2v_R^2\bar{v}_R^2]}{v_R^2 + \bar{v}_R^2},$$

$$M_{34} = \frac{2v_R\bar{v}_R(v_R^2 - \bar{v}_R^2)^2(2g_R^2 + 2g_V^2 + \lambda^2)}{v_R^2 + \bar{v}_R^2},$$

$$M_{35} = \frac{2\lambda[A_\lambda v_R\bar{v}_R + v_R^2(\lambda v_S + \mu_2) + \bar{v}_R^2(\lambda v_S + \mu_2) + v_R\bar{v}_R\mu_S]}{\sqrt{v_R^2 + \bar{v}_R^2}},$$

$$M_{44} = [8(g_R^2 + g_V^2)v_R^2\bar{v}_R^2 + (m_1^2 + m_2^2)(v_R^2 + \bar{v}_R^2) + \lambda^2(v_R^2 - \bar{v}_R^2)^2 \\ + 2(\lambda v_S + \mu_2)^2(v_R^2 + \bar{v}_R^2)] / (v_R^2 + \bar{v}_R^2),$$

$$M_{45} = -\frac{(v_R^2 - \bar{v}_R^2)\lambda(A_\lambda + \mu_S)}{\sqrt{v_R^2 + \bar{v}_R^2}},$$

$$M_{55} = m_S^2 + \lambda'^2(v_1^2 + v_2^2) + \lambda^2(v_R^2 + \bar{v}_R^2) + \mu_S^2 + 2\mu_S B_S.$$

( $\text{Re}\rho_1, \text{Re}\rho_2, \text{Re}\rho_3, \text{Re}\rho_4, \text{Re}S$ )

$$\rho_1 = \frac{v_1\phi_1^0 + v_2\phi_2^0}{\sqrt{v_1^2 + v_2^2}}, \quad \rho_2 = \frac{v_2\phi_1^0 - v_1\phi_2^0}{\sqrt{v_1^2 + v_2^2}}, \quad \rho_3 = \frac{v_R\delta^{c0} + \bar{v}_R\bar{\delta}^{c0}}{\sqrt{v_R^2 + \bar{v}_R^2}}, \quad \rho_4 = \frac{\bar{v}_R\delta^{c0} - v_R\bar{\delta}^{c0}}{\sqrt{v_R^2 + \bar{v}_R^2}}.$$

# Procedure adopted to derive light Higgs mass

- Choose  $\lambda'$ ,  $A_{\lambda'}$  and  $A_\lambda$  such that  $M_{13}$ ,  $M_{15}$  and  $M_{35}$  vanish. This would maximize the lightest Higgs boson mass.
- Lightest neutral scalar tree-level Higgs mass is found to be

$$M_{h_{tree}}^2 = 2M_W^2 \cos^2 2\beta + \lambda^2 v^2 \sin^2 2\beta, \quad (M_{h_{MSSM}}^2 = M_Z^2 \cos^2 2\beta)$$

- Using radiative corrections from top quark and stop squark

$$M_h^2 = (2M_W^2 \cos^2 2\beta + \lambda^2 \sin^2 2\beta)\Delta_1 + \Delta_2$$



where

$$\Delta_2 = \frac{3}{4\pi^2} \frac{m_t^4}{v^2} \left[ \frac{1}{2} \tilde{X}_t + t + \frac{1}{16\pi^2} \left( \frac{3}{2} \frac{m_t^2}{v^2} - 32\pi\alpha_3 \right) (\tilde{X}_t t + t^2) \right],$$

$$\Delta_1 = \left( 1 - \frac{3}{8\pi^2} \frac{m_t^2}{v^2} t \right), \quad t = \log \frac{M_S^2}{M_t^2}$$

$m_t$  is top running mass,  $\tilde{X}_t$  is the stop mixing,  $M_S$  is squark mass geometric mean and  $v = \sqrt{v_1^2 + v_2^2} \approx 174$  GeV.

# Higgs triplets and a singlet



- Green shaded region:  $X_t = 0$ , Blue shaded region:  $X_t = 6$ , Red shaded region: all possible values of parameters.
- Solid red region is for  $m_h$  between 124 GeV and 126 GeV.
- Solid black line: MSSM result with  $X_t = 6$  and  $M_S = 1.5$  TeV (or  $\tan\beta = 40$ ).
- Plot on the right panel shows that even for a relatively low stop mass and for small stop mixing the correct Higgs mass is produced.

# Pseudoscalar Higgs boson mass spectrum

Removing the two Goldstone states, elements of the  $3 \times 3$  matrix in a certain basis is

$$M_{11} = m_1^2 + m_2^2 + \lambda^2(v_R^2 + \bar{v}_R^2 + 2v_S^2) + 2\mu_2(2\lambda v_S + \mu_2),$$

$$M_{12} = \lambda\lambda' \sqrt{(v_1^2 + v_2^2)(v_R^2 + \bar{v}_R^2)},$$

$$M_{13} = \lambda(\mu_S - A_\lambda) \sqrt{v_R^2 + \bar{v}_R^2},$$

$$M_{22} = 2m_5^2 + \lambda'^2(v_1^2 + v_2^2 + 2v_S^2) + 2\mu(2\lambda'v_S + \mu),$$

$$M_{23} = \lambda'(2A_{\lambda'} - \mu_S) \sqrt{v_1^2 + v_2^2},$$

$$M_{33} = m_S^2 + \lambda^2(v_R^2 + \bar{v}_R^2) + \lambda'^2(v_1^2 + v_2^2) - \mu_S(2B_S - \mu_S).$$

Left-handed triplet Higgs fields decouple and give a complex mass-squared matrix

$$\begin{bmatrix} m_3^2 + \frac{g_L^2}{2}(v_1^2 - v_2^2) + g_V^2(-v_R^2 + \bar{v}_R^2) + (\lambda v_S + \mu_1)^2 & \lambda(M^2 - \lambda v_R \bar{v}_R + \lambda' v_1 v_2 - \mu_S v_S) - \lambda A_\lambda v_S - \mu_1 B_1 \\ \lambda(M^2 - \lambda v_R \bar{v}_R + \lambda' v_1 v_2 - \mu_S v_S) - \lambda A_\lambda v_S - \mu_1 B_1 & m_4^2 - \frac{g_L^2}{2}(v_1^2 - v_2^2) + g_V^2(v_R^2 - \bar{v}_R^2) + (\lambda v_S + \mu_1)^2 \end{bmatrix}$$

# Charged Higgs boson mass spectrum

Removing the two charged Goldstone states, elements of the  $2 \times 2$  matrix is

$$\begin{aligned}
 M_{11} &= -\frac{g_R^2(v_R^2 - \bar{v}_R^2)[v_1^4 + 2v_1^2(-v_2^2 + v_R^2 + \bar{v}_R^2) + v_2^4 + 2v_2^2(v_R^2 + \bar{v}_R^2)]}{(v_1^2 - v_2^2)(v_R^2 + \bar{v}_R^2)}, \\
 M_{12} &= \frac{2g_R^2 v_R \bar{v}_R \sqrt{v_1^4 + 2v_1^2(-v_2^2 + v_R^2 + \bar{v}_R^2) + v_2^2[v_2^2 + 2(v_R^2 + \bar{v}_R^2)]}}{v_R^2 + \bar{v}_R^2}, \\
 M_{22} &= [g_R^2 \{v_R^4(-v_1^2 + v_2^2 - 2v_R^2 - 2\bar{v}_R^2) + \bar{v}_R^4(-v_1^2 + v_2^2 + 2v_R^2 + 2\bar{v}_R^2) - 6(v_1^2 - v_2^2)v_R^2\bar{v}_R^2\} \\
 &\quad + 4g_V^2(v_R^2\bar{v}_R^4 - v_R^6 - v_R^4\bar{v}_R^2 + \bar{v}_R^6) - 2(m_1^2 - m_2^2)(v_R^2 + \bar{v}_R^2)^2] / (v_R^4 - \bar{v}_R^4).
 \end{aligned}$$

Left-handed charged triplet Higgs fields decouple and has a mass-squared matrix

$$\begin{pmatrix}
 g_V^2(\bar{v}_R^2 - v_R^2) + m_3^2 + \mu_1^2 & B_1\mu_1 \\
 B_1\mu_1 & g_V^2(v_R^2 - \bar{v}_R^2) + m_4^2 + \mu_1^2
 \end{pmatrix}$$

# Light doubly charged scalar

- Unlike the charged scalar, which is eaten up by  $W_R^\pm$ , one combination of doubly charged scalars from  $\delta^{c--}$  and  $\bar{\delta}^{c++}$  remains massless at tree level .
- This is due to the extra global symmetry of the model.
- The charge breaking vacuum turns out to be lower than the desired charge conserving vacuum.<sup>1</sup>
- Possible solutions with unbroken  $R$  Parity:
  - Planck scale corrections – Would require the right-handed symmetry breaking scale to be of order  $10^{11}$  GeV.<sup>2</sup>
  - Add new particles, eg:  $(1,1,3,0)$ .<sup>3</sup>
  - Stay minimal, but rely on radiative corrections to the Higgs mass.<sup>4,5</sup>

<sup>1</sup>R. Kuchimanchi, R. N. Mohapatra, Phys. Rev. D48, 4352 (1993) .

<sup>2,3</sup>C.S. Aulakh, K. Benakli, G. Senjanovic, Phys. Rev. Lett. 79, 2188 (1997);

C.S. Aulakh, A. Melfo, G. Senjanovic, Phys. Rev. D57, 4174 (1998).

<sup>3</sup> Z. Chacko, R. N. Mohapatra, Phys. Rev. D58, 015003 (1998).

<sup>4</sup> K.S. Babu, R. N. Mohapatra, Phys. Lett. B668, 404 (2008).

<sup>5</sup> K.S. Babu, A. Patra, to appear (2014).

# Doubly charged Higgs mass from radiative corrections

We have computed the full Majorana Yukawa contribution to the doubly charged Higgs mass.

The Higgs potential studied is a simplified version where electroweak VEVs are ignored.

$$V_F = \left| \lambda \text{Tr}(\Delta^c \bar{\Delta}^c) + \lambda'_{ab} \text{Tr}(\Phi_a^T \tau_2 \phi_b \tau_2) - M_R^2 \right|^2 + |\lambda|^2 |S|^2 \left| \text{Tr}(\Delta^c \Delta^{c\dagger}) + \text{Tr}(\bar{\Delta}^c \bar{\Delta}^{c\dagger}) \right|^2$$

$$V_{soft} = M_1^2 \text{Tr}(\Delta^c \Delta^{c\dagger}) + M_2^2 \text{Tr}(\bar{\Delta}^c \bar{\Delta}^{c\dagger}) + M_S^2 |S|^2 + \left\{ A_\lambda \lambda S \text{Tr}(\Delta^c \bar{\Delta}^c) - C_\lambda M_R^2 S + h.c. \right\}$$

$$V_D = \frac{g_R^2}{8} \sum \left| \text{Tr}(2\Delta^{c\dagger} \tau_a \Delta^c + 2\bar{\Delta}^{c\dagger} \tau_a \bar{\Delta}^c + \Phi_a \tau_a^T \Phi_a^\dagger) \right|^2 + \frac{g^2}{8} \sum_a \left| \text{Tr}(2\Delta^{c\dagger} \Delta^c + 2\bar{\Delta}^{c\dagger} \bar{\Delta}^c) \right|^2$$

- The charge-conserving VEV structure for the right-handed triplet Higgs boson is

$$\langle \Delta^c \rangle = \begin{bmatrix} 0 & v_R \\ 0 & 0 \end{bmatrix} \quad \langle \bar{\Delta}^c \rangle = \begin{bmatrix} 0 & 0 \\ \bar{v}_R & 0 \end{bmatrix}$$

while we can consider a charge-breaking vacuum given by

$$\langle \Delta^c \rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & v_R \\ v_R & 0 \end{bmatrix} \quad \langle \bar{\Delta}^c \rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & \bar{v}_R \\ \bar{v}_R & 0 \end{bmatrix}$$

- The  $SU(2)_R$  D-term vanishes for the charge-breaking vacuum.
- Charge-conserving vacuum has a positive D-term.
- The charge-breaking vacuum has a lower minima than the charge-conserving.
- The charge conserving vacuum leads to a massless (or negative squared mass) doubly charged scalar boson.

Right-handed doubly-charged Higgs boson mass-squared matrix (tree-level) is

$$M_{\delta^{++}}^2 = \begin{pmatrix} -2g_R^2(|v_R|^2 - |\bar{v}_R|^2) - \frac{\bar{v}_R}{v_R} Y & Y^* \\ Y & 2g_R^2(|v_R|^2 - |\bar{v}_R|^2) - \frac{v_R}{\bar{v}_R} Y \end{pmatrix}$$

where  $Y = \lambda A_\lambda S + |\lambda|^2 (v_R \bar{v}_R - \frac{M_R^2}{\lambda})$

Eigenvalues of the matrix

$$M_{\delta^{\pm\pm}}^2 = \frac{-Y(|v_R|^2 + |\bar{v}_R|^2) \pm \sqrt{(|v_R|^2 - |\bar{v}_R|^2)^2 |4g_R^2 v_R \bar{v}_R - Y|^2 + 4|v_R|^2 |\bar{v}_R|^2 |Y|^2}}{2|v_R| |\bar{v}_R|}$$

One of the eigenvalues is negative. Becomes zero if gauge couplings vanish.

However, radiative corrections can make the squared mass positive.

The problem solves itself through the one-loop Majorana Yukawa corrections, which break the accidental global symmetry of the tree-level Higgs potential.

- Corrections to pseudo-goldstone bosons must remain finite, which is a nontrivial check of the calculation.
- We neglect gauge couplings and calculate correction from Yukawa sector.
- This pseudo-Goldstone state is identified as


$$G^{++} = \frac{v_R^* \delta^{c--*} + \bar{v}_R \bar{\delta}^{c++}}{\sqrt{v_R^2 + \bar{v}_R^2}}$$

- The neutral Higgs boson couplings are written as


$$-\mathcal{L}_{\hat{X}} = P_i V_{ij} \hat{X}_j G^{++} G^{--} + Q_i V_{ij} \hat{X}_j n_1^2 + R_i V_{ij} \hat{X}_j n_2^2 + T_i V_{ij} \hat{X}_j \nu^c \nu^c$$

where  $\hat{X} \rightarrow$  Mass eigenstate,  $V \rightarrow$  Unitary diagonalizing matrix


# Feynman diagrams and corresponding contributions




$$M_1 = -\frac{i}{2} \left[ P^T M_h^{-2} Q \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2 - m_{n_1}^2} + P^T M_h^{-2} R \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2 - m_{n_2}^2} \right]$$




$$M_2 = 2i M_{\nu^c} P^T M_h^{-2} T \int \frac{d^4 k}{(2\pi)^4} \text{Tr} \left( \frac{\not{k} + M_{\nu^c}}{k^2 - M_{\nu^c}^2} \right)$$



$$M_3 = -\frac{i}{2} \frac{(f A_f v_R + f \lambda \bar{v}_R v_S)^2}{v_R^2 + \bar{v}_R^2} \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2 - m_{\tilde{e}^c}^2}$$



$$M_4 = -\frac{if^2v_R^2}{v_R^2 + \bar{v}_R^2} \int \frac{d^4k}{(2\pi)^4} \frac{1}{k^2}$$



$$M_5 = \frac{if^2v_R^2}{v_R^2 + \bar{v}_R^2} \int \frac{d^4k}{(2\pi)^4} \frac{1}{k^2 - m_{\tilde{e}^c}^2}$$

Adding all these contributions

- Quadratic divergences cancelled.
- Log divergences cancelled.

Final expression for one-loop correction to the mass of the right-handed doubly-charged Higgs boson:

$$\begin{aligned}
 M_{\delta^{\pm\pm}}^2 = & \frac{1}{16\pi^2} \frac{1}{v_R^2 + \bar{v}_R^2} \left[ f^2 v_R^2 m_{e^c}^2 \ln \left( \frac{m_{e^c}^2}{M_{\nu^c}^2} \right) + \frac{f^2 (\lambda \bar{v}_R v_S + A_f v_R)^2}{2(v_R^2 + \bar{v}_R^2)} \left\{ \ln \left( \frac{m_{e^c}^2}{M_{\nu^c}^2} \right) + 1 \right\} \right. \\
 & - \frac{f}{4} (\lambda \bar{v}_R v_S + 2f v_R^2 + A_f v_R) m_{\tilde{\nu}_1^c}^2 \ln \left( \frac{m_{\tilde{\nu}_1^c}^2}{M_{\nu^c}^2} \right) \\
 & \left. - \frac{f}{4} (-\lambda \bar{v}_R v_S + 2f v_R^2 - A_f v_R) m_{\tilde{\nu}_2^c}^2 \ln \left( \frac{m_{\tilde{\nu}_2^c}^2}{M_{\nu^c}^2} \right) \right]
 \end{aligned}$$

All the one-loop terms in the mass-square eigenvalues are of order  $M_{SUSY}^2 / 16\pi^2$ .

If right-handed gauge symmetry breaks at a high scale, the squared mass is negative. However, for  $v_R$  of order SUSY breaking scale, the mass squared is positive.

If SUSY is broken at the TeV scale, the doubly-charged Higgs boson mass has to be of electroweak symmetry breaking order ( $\sim 100$  GeV).

# Back to Light Neutral Higgs: Special Cases

## 1. Case without a Singlet Higgs $S$

- Higgs superpotential is:

$$W_{Higgs} = \mu_1 \text{Tr}(\Delta \bar{\Delta}) + \mu_2 \text{Tr}(\Delta^c \bar{\Delta}^c) + \frac{\mu}{2} \text{Tr}(\phi^T \tau_2 \phi \tau_2)$$

- Higgs mass is  $M_h = (M_Z^2 \cos^2 2\beta) \Delta_1 + \Delta_2$ .
- Same as in MSSM.

## 2. Case with the Singlet Higgs $S$ integrated out

- The Higgs superpotential is


$$W_{Higgs} = \mu_1 \text{Tr}(\Delta \bar{\Delta}) + \mu_2 \text{Tr}(\Delta^c \bar{\Delta}^c) + \frac{\mu}{2} \text{Tr}(\phi^T \tau_2 \phi \tau_2) + \varepsilon \text{Tr}(\Delta^c \bar{\Delta}^c)^2$$

- The lightest Higgs boson mass is


$$M_h = (2M_W^2 \cos^2 2\beta) \Delta_1 + \Delta_2.$$

# Higgs triplets and a heavy singlet

Maximum Higgs mass vs  $\tan\beta$



Maximum Higgs mass vs stop mass



- Green shaded region:  $X_t = 0$ , Blue shaded region:  $X_t = 6$ , Red shaded region: all possible values of parameters.
- Solid red region is for  $m_h$  between 124 GeV and 126 GeV.
- Solid black line: MSSM result with  $X_t = 6$  and  $M_S = 1.5$  TeV (or  $\tan\beta = 40$ ).
- Plot on the right panel shows that even for a relatively low stop mass and for small stop mixing the correct Higgs mass is produced.

# Symmetry breaking with Higgs doublets: Inverse Seesaw Models

Higgs sector is simple with doublets, and no triplets.

$$H_L(1, 2, 1, -1) = \begin{pmatrix} H_L^0 \\ H_L^- \end{pmatrix}, \quad \bar{H}_L(1, 2, 1, 1) = \begin{pmatrix} \bar{H}_L^+ \\ \bar{H}_L^0 \end{pmatrix}, \quad H_R(1, 1, 2, 1) = \begin{pmatrix} H_R^+ \\ H_R^0 \end{pmatrix},$$

$$\bar{H}_R(1, 1, 2, -1) = \begin{pmatrix} \bar{H}_R^0 \\ \bar{H}_R^- \end{pmatrix}, \quad \Phi_a(1, 2, 2, 0) = \begin{bmatrix} \phi_1^+ & \phi_2^0 \\ \phi_1^0 & \phi_2^- \end{bmatrix}_a, \quad (a = 1, 2)$$

- $H_R$  and  $\bar{H}_R$  breaks the right-handed symmetry.
- Allows right-handed symmetry breaking scale naturally of order TeV.
- $\Phi_a$  generates the quark and lepton masses and CKM mixing.
- Need extra singlet heavy neutrino  $N$  to generate small neutrino mass.

R. N. Mohapatra, J.W.F. Valle, Phys. Rev. D34, 1642 (1986).

- The non-zero vacuum expectation values of Higgs fields

$$\langle H_R^0 \rangle = v_R, \quad \langle \bar{H}_R^0 \rangle = \bar{v}_R, \quad \langle H_L^0 \rangle = v_L, \quad \langle \bar{H}_L^0 \rangle = \bar{v}_L, \quad \langle \phi_{1_a}^0 \rangle = v_{1_a}, \quad \langle \phi_{2_a}^0 \rangle = v_{2_a}$$

- Yukawa terms in the superpotential are

$$W_Y = \sum_{j=1}^2 Y_q^j Q^T \tau_2 \Phi_j \tau_2 Q^c + Y_l^j L^T \tau_2 \Phi_j \tau_2 L^c + if L^T \tau_2 \bar{H}_L N \\ + if^c L^{cT} \tau_2 \bar{H}_L N + \frac{\mu_N}{2} NN.$$

- Neutrino mass matrix – Inverse seesaw:

$$\begin{pmatrix} 0 & Y_l v_1 & f \bar{v}_L \\ Y_l v_1 & 0 & f^c \bar{v}_R \\ f \bar{v}_L & f^c \bar{v}_R & \mu_N \end{pmatrix}$$

S.M. Barr, Phys. Rev. Lett. 92, 101601 (2004).

- If  $\bar{v}_L \rightarrow 0$  and  $\mu_N \rightarrow 0$ , one of the eigenvalues of this matrix is zero.

- Consider the case with one bidoublet for simplicity.
- The Higgs only superpotential is


$$W_{Higgs} = i\mu_1 H_L^T \tau_2 \bar{H}_L + i\mu_1 H_R^T \tau_2 \bar{H}_R + \lambda \bar{H}_L^T \tau_2 \Phi \tau_2 \bar{H}_R \\ + \lambda H_L^T \tau_2 \Phi \tau_2 H_R + \mu \text{Tr} \left[ \Phi \tau_2 \Phi^T \tau_2 \right]$$

- Calculate the Higgs potential and the minimization conditions.
- Change the basis so that only one field gets EW VEV.
- Compute the neutral scalar Higgs boson mass-squared matrix.
- Mass of the lightest neutral Higgs boson


$$M_h = \left( 2M_W^2 \sin^4 \beta + \frac{M_W^4}{2M_W^2 - M_Z^2} \cos^4 \beta - \frac{M_W^2}{2} \sin^2 2\beta + \lambda^2 v^2 \sin^2 2\beta \right) \Delta_1 + \Delta_2$$

# Higgs doublets: Inverse seesaw models

Maximum Higgs mass vs  $\tan\beta$



Maximum Higgs mass vs stop mass



- Green shaded region:  $X_t = 0$ , Blue shaded region:  $X_t = 6$ , Red shaded region: all possible values of parameters.
- Solid red region is for  $m_h$  between 124 GeV and 126 GeV.
- Solid black line: MSSM result with  $X_t = 6$  and  $M_S = 1.5$  TeV (or  $\tan\beta = 40$ ).
- Plot on the right panel shows that even for a relatively low stop mass and for small stop mixing the correct Higgs mass is produced.

# Universal Seesaw Models

Higgs sector of the model is

$$H_L(1,2,1,-1) = \begin{pmatrix} H_L^0 \\ H_L^- \end{pmatrix}, \quad \bar{H}_L(1,2,1,1) = \begin{pmatrix} \bar{H}_L^+ \\ \bar{H}_L^0 \end{pmatrix}, \quad H_R(1,1,2,1) = \begin{pmatrix} H_R^+ \\ H_R^0 \end{pmatrix},$$

$$\bar{H}_R(1,1,2,-1) = \begin{pmatrix} \bar{H}_R^0 \\ \bar{H}_R^- \end{pmatrix}, \quad S(1,1,1,0)$$

- $H_R$  and  $\bar{H}_R$  breaks the right-handed symmetry.
- No bidoublet to generate the quarks and lepton masses and mixings.
- Need extra heavy quarks and leptons

$$P(3,1,1,-\frac{4}{3}), \quad R(3,1,1,\frac{2}{3}), \quad E(1,1,1,2)$$

$$P^c(3,1,1,\frac{4}{3}), \quad R^c(3,1,1,-\frac{2}{3}), \quad E^c(1,1,1,-2)$$

A. Davidson, KC. Wali, Phys. Rev. Lett. 59, 393 (1987);  
K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. 62, 1079 (1989).

- An optional heavy singlet neutrino  $N$ .

- Yukawa coupling terms in the superpotential

$$\begin{aligned}
W_Y &= Y_u Q \bar{H}_L P - Y_d Q H_L R - Y_l L H_L E + Y_\nu L \bar{H}_L N \\
&+ Y_u^c Q^c \bar{H}_R P^c - Y_d^c Q^c H_R R^c - Y_l^c L^c H_R E^c + Y_\nu^c L^c \bar{H}_R N \\
&+ m_u P P^c + m_d R R^c + m_l E E^c + m_\nu N N
\end{aligned}$$

- Mass of fermions are generated via the seesaw mechanism.

$$M_u = \begin{pmatrix} 0 & Y_u \bar{\nu}_L \\ Y_u^c \bar{\nu}_R & m_u \end{pmatrix}$$

- Neutrino mass can also be generated at the two-loop level from  $W_L$  and  $W_R$  exchange.



- The Higgs only superpotential is

$$W_{Higgs} = S \left( i\lambda H_L^T \tau_2 \bar{H}_L + i\lambda H_R^T \tau_2 \bar{H}_R - M^2 \right).$$

- Mass of the lightest neutral scalar Higgs boson

$$M_h = \left( \frac{M_W^4}{2M_W^2 - M_Z^2} \cos^2 2\beta + \lambda^2 v^2 \sin^2 2\beta \right) \Delta_1 + \Delta_2$$

# Higgs doublets: Universal seesaw



- Green shaded region:  $X_t = 0$ , Blue shaded region:  $X_t = 6$ , Red shaded region: all possible values of parameters.
- Solid red region is for  $m_h$  between 124 GeV and 126 GeV.
- Solid black line: MSSM result with  $X_t = 6$  and  $M_S = 1.5$  TeV (or  $\tan\beta = 40$ ).
- Plot on the right panel shows that even for a relatively low stop mass and for small stop mixing the correct Higgs mass is produced.

# Universal seesaw model without singlet

- Higgs superpotential terms are

$$W_{Higgs} = i\mu_1 H_L^T \tau_2 \bar{H}_L + i\mu_1 H_R^T \tau_2 \bar{H}_R.$$

- Calculate the Higgs potential and the minimization conditions.
- Calculate the Higgs boson mass-squared matrix.
- Lightest neutral scalar Higgs boson mass

$$M_h = \left( M_Z^2 \cos^2 2\beta \right) \Delta_1 + \Delta_2.$$

- Same result as in MSSM.

# $E_6$ motivated left-right SUSY model

- Low energy manifestation of superstring theory.
- Matter multiplets belong to 27 of  $E_6$  group.
- Previously discussed by others but some parameters were zero, here we keep everything.
- Particle spectrum given as

$$X^c(\bar{3}, 1, 2, -\frac{1}{6}) = \begin{pmatrix} h^c & u^c \end{pmatrix}_L, \quad Q(3, 2, 1, \frac{1}{6}) = \begin{pmatrix} u & d \end{pmatrix}_L, \quad h(3, 1, 1, -\frac{1}{3}) = h_L,$$

$$L^c(1, 1, 2, \frac{1}{2}) = \begin{pmatrix} e^c & n \end{pmatrix}_L, \quad E(1, 2, 1, -\frac{1}{2}) = \begin{pmatrix} \nu_E & E \end{pmatrix}_L, \quad d^c(\bar{3}, 1, 1, \frac{1}{3}) = d_L^c,$$

$$F(1, 2, 2, 0) = \begin{pmatrix} \nu_e & E^c \\ e & N_E^c \end{pmatrix}_L, \quad N^c(1, 1, 1, 0) = N_L^c.$$

- Discrete R-parity symmetry under which

$$(u, d, e, \nu_e) \in \text{Even}, \quad (h, E, n, N_E^c, \nu_E) \in \text{Odd}$$

- The Higgs fields are identified as

$$H_L(1,2,1,-1) = \begin{pmatrix} H_L^0 \\ H_L^- \end{pmatrix} = \begin{pmatrix} \tilde{\nu}_E \\ \tilde{E} \end{pmatrix}, \quad H_R(1,1,2,1) = \begin{pmatrix} H_R^+ \\ H_R^0 \end{pmatrix} = \begin{pmatrix} \tilde{e}^c \\ \tilde{n} \end{pmatrix},$$

$$\Phi(1,2,2,0) = \begin{pmatrix} \phi_1^+ & \phi_2^0 \\ \phi_1^0 & \phi_2^- \end{pmatrix} = \begin{pmatrix} \tilde{E}^c & \tilde{N}_E^c \\ \tilde{\nu}_e & \tilde{e} \end{pmatrix}.$$

- The superpotential is

$$W = \lambda_1 Q d^c E + \lambda_2 Q X^c F + \lambda_3 h X^c L^c + \lambda_4 F L^c E + \lambda_5 F N^c F + \lambda_6 h d^c N^c$$

- The quark and lepton masses are generated from this superpotential.
- The neutrino mass matrix is a  $3 \times 3$  matrix in basis  $(\nu_E, N_E^c, n)$  given as

$$\begin{pmatrix} 0 & \lambda_4 \langle \tilde{n} \rangle & \lambda_4 \langle \tilde{N}_E^c \rangle \\ \lambda_4 \langle \tilde{n} \rangle & 0 & \lambda_4 \langle \tilde{\nu}_E \rangle \\ \lambda_4 \langle \tilde{N}_E^c \rangle & \lambda_4 \langle \tilde{\nu}_E \rangle & 0 \end{pmatrix}$$

- The Higgs only superpotential is

$$W_{Higgs} = \lambda H_L^T \tau_2 \Phi \tau_2 H_R + \mu \text{Tr} \left[ \Phi^T \tau_2 \Phi \tau_2 \right].$$


- The non-zero vacuum expectation values are given as

$$\langle H_R^0 \rangle = v_R, \quad \langle H_L^0 \rangle = v_L, \quad \langle \phi_1^0 \rangle = v_1, \quad \langle \phi_2^0 \rangle = v_2$$

- Calculate the Higgs potential and the minimization conditions.
- Change basis so that only one field gets EW vev.
- Compute the Higgs boson mass-squared matrix.
- Mass of the lightest neutral Higgs boson

$$M_h = \left( 2M_W^2 \cos^2 2\beta \right) \Delta_1 + \Delta_2$$

# Experimental search for doubly-charged Higgs boson at LHC



Direct pair-production of doubly-charged Higgs Boson


The mass limit on the right-handed doubly-charged Higgs boson from ATLAS experiment.

| BR( $H_R^{\pm\pm} \rightarrow \ell^\pm \ell'^\pm$ ) | 95% CL lower limit on $m(H_R^{\pm\pm})$ [GeV] |      |                   |      |                 |      |
|---|---|------|-------------------|------|-----------------|------|
|   | $e^\pm e^\pm$                                 |      | $\mu^\pm \mu^\pm$ |      | $e^\pm \mu^\pm$ |      |
|   | exp.  | obs. | exp.              | obs. | exp.            | obs. |
| 100%  | 329   | 322  | 335               | 306  | 303             | 310  |
| 33%   | 241   | 214  | 247               | 222  | 220             | 195  |
| 22%   | 203   | 199  | 223               | 212  | 194             | 187  |
| 11%   | 160   | 151  | 184               | 176  | 153             | 151  |

S. Chatrchyan et al. [CMS Collaboration], Eur. Phys. J. C 72, 2189 (2012);  
 G. Aad et al. [ATLAS Collaboration], Eur.Phys. J. C 72, 2244 (2012).

# New signals of doubly-charged particles at the LHC

Production and decay of the doubly-charged Higgsino.




The final state signal is 4 leptons and missing energy (a new channel for doubly-charged Higgs boson search at the LHC).

Three cases we must consider


1.  $p p \rightarrow \delta_R^{++} \delta_R^{--} \rightarrow l^+ l^+ l^- l^-$
2.  $p p \rightarrow \tilde{\delta}_R^{++} \tilde{\delta}_R^{--} \rightarrow \delta_R^{++} \delta_R^{--} \tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow l^+ l^+ l^- l^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$
3.  $p p \rightarrow \tilde{\delta}_L^{++} \tilde{\delta}_L^{--} \rightarrow \tilde{l}^+ l^+ \tilde{l}^- l^- \rightarrow l^+ l^+ l^- l^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$

We take two parameter space

- BP1 for which  $M_{\delta_R^{++}} = 300$  GeV,  $M_{\tilde{\delta}_R^{++}} = 500$  GeV and  $M_{\tilde{\chi}_1^0} = 80$  GeV .
- BP2 for which  $M_{\delta_R^{++}} = 300$  GeV,  $M_{\tilde{\delta}_R^{++}} = 400$  GeV and  $M_{\tilde{\chi}_1^0} = 80$  GeV .



Direct production of the light doubly-charged Higgsinos and Higgs boson at LHC at 14 TeV.




$\Delta R$  plot for the same-sign and opposite-sign final state leptons.

The same-sign lepton plot peaks at a low value of  $\Delta R$  while the opposite-sign leptons peak at a much higher value.

Expected as the same-sign leptons come from the decay of one particle while the opposite sign leptons come from much further apart.

Measurement at the LHC for a four lepton final state can give definite indication of existence of doubly-charged particles if distribution is similar to our analysis.



Invariant Mass plot for the same- sign and opposite-sign final state leptons.

No events between 80 GeV and 100 GeV in the same-sign lepton plot because of the Z peak cut applied.

A clear peak in the same sign invariant mass while no such peak in the opposite sign plot at a mass of 300 GeV.

The Peak came due to the decay of the right-handed doubly-charged Higgs bosons into two same-sign leptons.

Difficult to see in experiments without a priori knowledge of the Higgs boson mass.

If seen, will be a definite signature of a doubly-charged particle.

# Conclusions



- ❧ The Left-Right Supersymmetric models solve many of the problems in the Standard Model.
- ❧ Examined various models with different symmetry breaking sectors.
- ❧ The tree-level neutral Higgs boson mass can be significantly increased.
- ❧ Experimentally observed Higgs boson mass of 125 GeV can be easily achieved with low stop mass and mixing.
- ❧ Models with triplets admit zero mass states of doubly-charged Higgs bosons which remains light after radiative corrections.

THANK YOU

## The kinematic cuts applied

- The final state lepton must have a rapidity cut  $-2.5 < \eta_l < 2.5$ .
- $\Delta R_{ll} > 0.2$  between the final state leptons  
where  $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$  and  $\phi$  is the azimuthal angle.
- Each lepton must have a transverse momentum  $p_T > 15$  GeV.
- Invariant mass cut between the opposite sign same flavor leptons such that  $M_{inv}^{OS} > 10$  GeV.
- A further invariant mass cut of  $80$  GeV  $> M_{inv}^{OS} > 100$  GeV to get rid of the Z-boson peak.

We study the MET,  $\Delta R$  and the invariant mass of the final state leptons.